

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
1 April 2004 (01.04.2004)

PCT

(10) International Publication Number  
WO 2004/026224 A2

- (51) International Patent Classification<sup>7</sup>: A61K (74) Agent: CHOI, Kyu Pal; 4th Floor, Halla Classic Building, 824-11, Yeoksam-dong, Kangnam-gu, Seoul 135-080 (KR).
- (21) International Application Number: PCT/KR2003/001902
- (22) International Filing Date: 18 September 2003 (18.09.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
10-2002-0056836  
18 September 2002 (18.09.2002) KR
- (71) Applicant (for all designated States except US): YON-SUNG FINE CHEMICAL CO. LTD. [KR/KR]; 51-301, Science Technology Park, Sungkyunkwan University, 300, Chonchon-dong, Changan-gu, Suwon, Kyonggi-do 440-746 (KR).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): KIM, Yong Hyun [KR/KR]; 73-73, Hwagok 5-dong, Gangseo-gu, Seoul 157-015 (KR). LEE, Yiu Suk [KR/KR]; Woosung Apt. #304-407, 105, Sadang-dong, Dongjak-gu, Seoul 156-090 (KR).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR PREPARING PROSTAGLANDIN DERIVATIVES AND STARTING MATERIALS FOR THE SAME

(57) Abstract: The present invention provides a process for preparing benzoprostacycline derivatives of formula (I), i.e. 5,6,7-trinor-4,8-inter-m-phenylene PGI<sub>2</sub> derivatives, and vinyl tin compounds of formula (III) as starting materials for the same.

BEST AVAILABLE COPY


WO 2004/026224 A2

PROCESS FOR PREPARING PROSTAGLANDIN DERIVATIVES AND  
 STARTING MATERIALS FOR THE SAME

TECHNICAL FIELD

5

The present invention relates to a process for preparing benzoprostacycline derivatives of the following formula:




, wherein  $R^1$  represents a cation, H, or  $C_{1-12}$ alkyl,

10  $R^2$  and  $R^3$  each represent H or a hydroxy protective group, preferably, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, phenyldimethylsilyl, or tetrahydropyranyl,

$R^4$  represents H or  $C_{1-3}$ alkyl, and

$R^5$  represents H or  $C_{1-6}$ alkyl,

15 i.e. 5,6,7-trinor-4,8-inter-m-phenylene  $PGI_2$  derivatives. The present invention also relates to vinyl tin compounds of the following formula:




, wherein  $R^{3a}$  represents a hydroxy protective group, preferably, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, phenyldimethylsilyl, or  
 20 tetrahydropyranyl, and

$R^4$  and  $R^5$  are each as defined above (Bu: Butyl),

as starting materials for the above process.

## BACKGROUND ART

5 Benzoprostacycline derivatives of formula (I) are known in the United States Patent No. 5,202,447 and Tetrahedron Lett. 31, 4493 (1990), and their general synthetic method is depicted in the following reaction scheme:



(Et: Ethyl, Ts: Tosyl, THF: Tetrahydrofuran, LAH: Lithium aluminum hydride,  
 10 TBS: t-Butyldimethylsilyl, DMF: Dimethylformamide, pyr.: Pyridine, Ac: Acetyl, Me: Methyl)

Specifically, the above method comprises the steps of: i) forming a diol compound from a benzofuran ring via Prins reaction; ii) introducing a substituent to the benzene ring via Grignard reaction; iii) selectively protecting the diol, and then, introducing an  $\omega$ -chain thereto via Wittig reaction; and iv) performing reduction, deacetylation and ester hydrolysis to the obtained compound under Luche condition. However, the method has drawbacks in that the diastereomeric mixture formed from reduction of  $\text{C}_{15}$  ketone after  $\omega$ -

15

chain introduction has extremely low stereo-selectivity, i.e. the isomers ratio of 45:55, and a very low yield, i.e. 40%, of the desired isomer.


## DISCLOSURE OF THE INVENTION

5

The present inventors have performed extensive studies to develop a process for preparing the compound of formula (I) which is not only economic but also efficient by solving the above-described problems of the known processes in the art. As a result, the present inventors invented novel compounds containing an  $\omega$ -chain, and found that the above purpose could be attained by using the first step of performing 1,4-addition reaction of the  $\omega$ -chain-containing compounds to  $\alpha,\beta$ -unsaturated ketones to prepare the compound of formula (I). Thus, the present invention was completed.

One aspect of the present invention provides a process for preparing the compound of the following formula:

15



(I)

, wherein  $\text{R}^1$  represents a cation, H, or  $\text{C}_{1-12}$ alkyl,


$\text{R}^2$  and  $\text{R}^3$  each represent H or a hydroxy protective group, preferably, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, phenyldimethylsilyl or tetrahydropyranyl,

20

$\text{R}^4$  represents H or  $\text{C}_{1-3}$ alkyl, and

$\text{R}^5$  represents H or  $\text{C}_{1-6}$ alkyl, comprising the steps of:


(1) converting the compound of the following formula:



, wherein  $R^{3a}$  represents a hydroxy protective group, preferably, trimethylsilyl, triethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl, phenyldimethylsilyl, or tetrahydropyranyl, and

$R^4$  and  $R^5$  are each as defined above (Bu: Butyl),

into its cuprate, and then, performing stereo-specific 1,4-addition reaction of the cuprate to an  $\alpha,\beta$ -unsaturated ketone of the following formula:




, wherein  $R^1$  is as defined above,

$R^{2a}$  is as defined above for  $R^{3a}$ , and


$R'$  represents a hydroxy protective group, preferably, methyl, methoxymethyl, methoxyethyl, or a substituted methyl ether such as benzyloxymethyl, p-benzyloxymethyl, etc.,

to form a compound of the following formula (IV);




, wherein  $R^1$ ,  $R^{2a}$ ,  $R^{3a}$ ,  $R^4$ ,  $R^5$  and  $R'$  are each as defined above;

(2) reducing the ketone of cyclopentanone in the compound of formula (IV) to form an  $\alpha$ -alcohol compound of the following formula:



, wherein R<sup>1</sup>, R<sup>2a</sup>, R<sup>3a</sup>, R<sup>4</sup>, R<sup>5</sup> and R' are each as defined above;


(3) substituting the  $\alpha$ -alcohol in the compound of formula (V) with a halide to form a  $\beta$ -halide compound of the following formula:



, wherein R<sup>1</sup>, R<sup>2a</sup>, R<sup>3a</sup>, R<sup>4</sup>, R<sup>5</sup> and R' are each as defined above, and

X represents halo, preferably, chloro;


(4) deprotecting a hydroxy protective group of the compound of formula (VI) to form a compound of the following formula:



, wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and X are each as defined above;

(5) performing intramolecular cyclization to the compound of formula (VII) to form the compound of formula (I).

Another aspect of the present invention provides a compound of the following formula:




, wherein  $R^{3a}$ ,  $R^4$  and  $R^5$  are each as defined above.

Hereinafter, the present invention will be described in detail.

5

The compound of formula (III) used as a starting material for the preparation process of the present invention is novel, and the examples of its preparation processes include one comprising the step of reacting a compound of the following formula:



10

with a compound of the following formula:




, wherein  $R^4$  and  $R^5$  are each as defined above,

, and then protecting the resulting hydroxy group (process (a)); and

one comprising the step of reacting the compound of formula (VIII) with a

15

compound of the following formula:



, wherein  $R^4$  and  $R^5$  are each as defined above (process (b)).

#### a) Process (a)


20

The compound of formula (VIII) is dissolved in THF, and then, the resulting solution is cooled to  $-78\text{ }^{\circ}\text{C}$ .  $n\text{-BuLi}$  is slowly added thereto, and the mixture is stirred at the above temperature for 1 hour. Then, the compound of formula (IX) diluted in THF is added thereto, and the mixture is stirred for 5 minutes. Then, a hydroxy protective agent (e.g. TBSCl) is added thereto to obtain the compound of formula (III).

## b) Process (b)

The compound of formula (VIII) is dissolved in THF, and then, the resulting solution is cooled to  $-78^{\circ}\text{C}$ .  $n\text{-BuLi}$  is slowly added thereto, and the mixture is stirred at the above temperature for 1 hour. Then, the compound of formula (X) diluted in THF is added thereto, and the mixture is stirred for 5 minutes to stop the reaction with an aqueous solution of  $\text{NH}_4\text{Cl}$ . The reaction solution is extracted with diethyl ether, and the extract is concentrated. Thereto are added methanol and cerium chloride in the order. The ketone is reduced with sodium borohydride, and brine is added thereto. The mixture is extracted with hexane and the extract is concentrated. The concentrate is dissolved in DMF and a hydroxy protective agent (e.g. TBSCl) is added thereto to obtain the compound of formula (III).

The preparation process according to the present invention is depicted in the following reaction scheme:



Each step of the preparation process according to the present invention will be explained below.



(1) First step: Preparation of the compound of formula (IV) from the compound of formula (II) and the compound of formula (III)

5       The compound of formula (IV) is prepared by conducting 1,4-addition reaction of a cuprate formed from a vinyl tin of formula (III) to an  $\alpha,\beta$ -unsaturated ketone in the compound of formula (II). In this reaction, the cuprate is added to the opposite side of the alkoxy group ( $-\text{OR}^{2a}$ ) substituted in the cyclopentanone ring due to steric hindrance from the alkoxy group, and so has trans-configuration with the alkoxy group. The added  $\omega$ -  
10 chain causes another steric hindrance to the  $\alpha$ -chain of the cyclopentanone to give the compound of formula (IV) with the stereochemistry of trans configuration between  $\omega$ -chain and  $\alpha$ -chain.

(2) Second step: Preparation of the compound of formula (V) by reduction of the  
15 compound of formula (IV)

The reduction of ketone present in the compound of formula (IV) may be carried out by using various kinds of metal hydride. The examples of metal hydride which may be used include sodium borohydride ( $\text{NaBH}_4$ ), L-selectride, N-selectride, or K-selectride.  
20 L-selectride is preferable because a bulkier metal hydride has greater steric hindrance from the alkoxy group ( $\text{R}^{2a}\text{O}-$  or  $\text{R}^{3a}\text{O}-$ ) substituted in the compound of formula (IV), and so the hydride attacks the cyclopentanone ring on the opposite side of the alkoxy group so that the desired  $\alpha$ -alcohol can be selectively obtained. For this reaction, 1 to 3 equivalents of L-selectride is added to 1 equivalent of the compound of formula (IV) dissolved in a  
25 solvent at  $-78^\circ\text{C}$ , and the mixture is stirred for 1 to 2 hours to stop the reaction with 30%  $\text{H}_2\text{O}_2$ . The reaction solution is stirred at  $0^\circ\text{C}$  for 30 minutes, extracted with diethyl ether, and concentrated to stereo-selectively obtain the compound of formula (V). The compound is used in the subsequent reaction without any further purification. The examples of solvent which may be used include tetrahydrofuran, diethyl ether, and  
30 dichloromethane.

(3) Third step: Preparation of the compound of formula (VI) by halide substitution of the compound of formula (V)

5           The  $\alpha$ -alcohol of formula (V) is substituted with a halide with various routes. In this process, the alcohol is converted into a leaving group, and then, the leaving group is converted into the  $\beta$ -halide via  $S_N2$  reaction where the halide acts as a nucleophile. The examples of the leaving group include p-toluene sulfonate, methane sulfonate, trifluoromethane sulfonate, and the like, and the examples of the halide nucleophile include  
10   lithium halide, tetrabutylammonium halide, and the like. A method to directly convert the alcohol into the halide using trialkylphosphine and carbon tetrahalide may also be used. Preferable examples of the trialkylphosphine include tri-n-butylphosphine, triphenylphosphine and the like.

15           (4) Fourth step: Preparation of the compound of formula (VII) by deprotecting the compound of formula (VI)

          The benzene ring present in the compound of formula (VI), and optionally, the hydroxy protective group(s) of cyclopentanone ring and/or of  $\omega$ -chain can be deprotected  
20   under an acidic condition. The examples of acid which can be used include diluted hydrochloric acid, p-toluenesulfonic acid, pyridinium p-toluenesulfonic acid and the like, and the examples of solvent which can be used include methanol, ethanol, tetrahydrofuran, dichloromethane, acetone and mixed solvents thereof, etc.

25           (5) Fifth step: Preparation of the compound of formula (I) by intramolecular cyclization of the compound of formula (VII)

          The compound of formula (I) can be prepared from the compound of formula (VII) via  $S_N2$  reaction of a phenolic hydroxy group with a  $\beta$ -halide in the presence of  
30   various kinds of base. In this reaction, the examples of base include conventional base,

for example, trialkylamine such as trimethylamine, triethylamine or diisopropylethylamine, and an inorganic base such as NaH, K<sub>2</sub>CO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub>, KHCO<sub>3</sub> or NaHCO<sub>3</sub>, particularly, K<sub>2</sub>CO<sub>3</sub>. The solvent may be polar or non-polar if it does not affect the reaction, but dimethylformamide (DMF), dimethylacetamide (DMA), tetrahydrofuran (THF),  
5 acetonitrile (CH<sub>3</sub>CN), and so on are preferable, and CH<sub>3</sub>CN is particularly preferable. An appropriate reaction temperature is about 0 to 100 °C.

### BEST MODE FOR CARRYING OUT THE INVENTION

10 The present invention will be more specifically illustrated by the following examples. However, the following examples should not be construed as limiting the scope of the present invention in any way.


**Preparation: Synthesis of t-butyl dimethyl(2-methyl-1-(2-tributylstanyl-vinyl)-4-hexynyloxy)silane**  
15

Trans-bis(tributyl)tin ethylene (110 g, 181 mmol) was added to THF (500 ml), and the mixture was cooled to -78 °C. BuLi (72 ml, 2.5 M in hexane, 181 mmol) was added thereto dropwise, and the mixture was stirred at the above temperature for 1 hour. Then,  
20 2-methylhexyn-4-ol methoxymethylamide (25.5 g, 151 mmol) dissolved in THF (30 ml) was added thereto. The resulting mixture was stirred at the above temperature for 15 minutes, and then, a saturated aqueous solution of NH<sub>4</sub>Cl (100 ml) was added thereto. The reaction temperature was elevated to room temperature, and then, the reaction mixture was extracted with hexane. The obtained organic layer was dried over anhydrous  
25 magnesium sulfate, and then, filtered off, and concentrated under reduced pressure. The residue was dissolved in 1 l of methanol, and then, CeCl<sub>3</sub> (50 g, 136 mmol) was added thereto. The mixture was cooled to 0 °C, and then, sodium borohydride (5.2 g, 136 mmol) was added thereto and the resulting mixture was stirred for 10 minutes. Brine of 500 ml was added thereto, and the mixture was extracted with hexane. The organic layer  
30 was dried over anhydrous magnesium sulfate, and then, filtered off, and concentrated under

reduced pressure. Thereto were added DMF (500 ml), imidazole (27 g, 400 mmol), and TBSCl (30 g, 200 mmol) in the order, and the mixture was stirred for 12 hours. Upon completion of the reaction, the reaction solution was extracted with hexane, and the organic layer was dried over anhydrous magnesium sulfate, and then, filtered off, and concentrated under reduced pressure. The concentrate was subjected to column chromatography on a silica gel to obtain colorless transparent t-butyldimethyl(2-methyl-1-(2-tributylstanyl-vinyl)-4-hexynyloxy)silane (60 g, yield: 74%).

<sup>1</sup>H NMR: 0.89 (m, 28H), 1.30-1.58 (m, 14H), 1.79 (t, 3H), 1.92-2.21 (m, 2H), 3.89 (dd, 1/2H), 4.06 (dd, 1/2H), 5.82-6.08 (m, 2H)

**Example 1:** Synthesis of 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-oxo-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester



CuCN (1.21 g, 13.5 mmol) was added to THF (15 ml), and the mixture was cooled to 0 °C. MeLi (20.6 ml, 1.44 M in Et<sub>2</sub>O, 29.7 mmol) was added thereto dropwise. t-Butyldimethyl(2-methyl-1-(2-tributylstanyl-vinyl)-4-hexynyloxy)silane (8.2 g, 15.2 mmol) was added thereto dropwise, and the mixture was stirred at room temperature for 2 hours. The reaction solution was cooled to -65 °C, and thereto was added 4-{3-[3-(tert-butyl-dimethyl-silanyloxy)-5-oxo-cyclopent-1-enyl]-2-methoxymethoxy-phenyl}-butyric acid methyl ester (4.3 g, 9.6 mmol) dissolved in 15 ml of THF. The reaction temperature was elevated to -35 °C. The mixture was stirred for 5 minutes, and then, added to a solution of NH<sub>4</sub>Cl-NH<sub>3</sub> (9:1). The resulting mixture was stirred for 15 minutes, and


extracted with diethyl ether. The organic layer was dried over anhydrous magnesium sulfate, and filtered off, and concentrated under reduced pressure. The concentrate was subjected to column chromatography on a silica gel (hexane:ethyl acetate=8:1) to obtain 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-oxo-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester (5.6 g, yield: 83%).

**Example 2:** Synthesis of 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-hydroxy-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester



4-(3-{3-(tert-Butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-oxo-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester of 2.6 g (3.67 mmol) was dissolved in THF (40 ml). The solution was cooled to -78 °C, and then, L-selectride (1 M THF solution, 7.35 mmol) was slowly added thereto dropwise. The mixture was stirred at the above temperature for 2 hours, and the reaction was stopped with hydrogen peroxide (30% aqueous solution, 16 mmol). The reaction solution was stirred at 0 °C for 30 minutes, and then, extracted with diethyl ether. The combined organic layer was washed with water and brine, and then, dried over magnesium sulfate, filtered off, and concentrated under reduced pressure. The concentrate was subjected to column chromatography on a silica gel (hexane:ethyl acetate=4:1) to obtain 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-hydroxy-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester (2.3 g, yield: 90%).

**Example 3:** Synthesis of 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-chloro-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester



A mixture of 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-hydroxy-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester of 506 mg (0.72 mmol), triphenyl phosphine (350 mg, 1.34 mmol), CH<sub>3</sub>CN (6 ml), pyridine (112 mg, 1.42 mmol), and CCl<sub>4</sub> (240 mg, 1.55 mmol) was stirred for 12 hours. The reaction solution was concentrated under reduced pressure, and then, subjected to column chromatography on a silica gel to obtain 4-(3-{3-(tert-butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-chloro-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester (426 mg, yield: 82%).


**Example 4:** Synthesis of 4-{3-[5-chloro-3-hydroxy-2-(3-(hydroxy-4-methyl-oct-1-en-6-ynyl)-cyclopentyl)-2-hydroxy-phenyl]-butyric acid methyl ester



4-(3-{3-(tert-Butyl-dimethyl-silanyloxy)-2-[3-(tert-butyl-dimethyl-silanyloxy)-4-methyl-oct-1-en-6-ynyl]-5-chloro-cyclopentyl}-2-methoxymethoxy-phenyl)-butyric acid methyl ester of 939 mg (1.3 mmol) was dissolved in a mixed solution of

methanol/dichloromethane (5:2) of 14 ml, and p-toluenesulfonic acid (895 mg, 5.2 mmol) was added thereto. The mixture was stirred at room temperature for 12 hours. Upon completion of the reaction, the reaction mixture was washed with an aqueous solution of sodium bicarbonate, and then, dried over anhydrous magnesium sulfate, filtered off, and concentrated under reduced pressure. The concentrate was subjected to column chromatography on a silica gel to obtain 4-{3-[5-chloro-3-hydroxy-2-(3-(hydroxy-4-methyl-oct-1-en-6-ynyl)-cyclopentyl)-2-hydroxy-phenyl]-butyric acid methyl ester (304 mg, yield: 52%).

**Example 5:** Synthesis of 4-[2-hydroxy-3-(3-hydroxy-4-methyl-oct-1-en-6-ynyl)-2,3,3a,8a-tetrahydro-1H-8-oxa-cyclopenta[a]inden-7-yl]-butyric acid methyl ester



4-{3-[5-Chloro-3-hydroxy-2-(3-(hydroxy-4-methyl-oct-1-en-6-ynyl)-cyclopentyl)-2-hydroxy-phenyl]-butyric acid methyl ester of 1.12 g (2.5 mmol) was dissolved in acetonitrile of 10 ml. Then, potassium carbonate of 691 mg (5.0 mmol) was added thereto, and the mixture was stirred at room temperature for 4 hours. Upon completion of the reaction, the reaction solution was concentrated under reduced pressure, and thereto was added water of 10 ml. The resulting mixture was extracted three times with ethyl acetate of 10 ml. The combined organic layer was dried over anhydrous magnesium sulfate, and then, filtered off, and concentrated under reduced pressure. The concentrate was subjected to column chromatography on a silica gel (hexane:ethyl acetate=1:2) to obtain 4-[2-hydroxy-3-(3-hydroxy-4-methyl-oct-1-en-6-ynyl)-2,3,3a,8a-tetrahydro-1H-8-oxa-cyclopenta[a]inden-7-yl]-butyric acid methyl ester (980 mg, yield: 95%).


### INDUSTRIAL APPLICABILITY

The process of the present invention can prevent yield decrease caused by the  
5 formation of isomers in  $\omega$ -chain reduction, and by the separation of isomers in the final  
step. According to the present process, the desired product, the compound of formula (I),  
can be readily purified and obtained in a highly stereo-specific manner, and so prepared  
economically and efficiently.




## WHAT IS CLAIMED IS:

1. A process for preparing a compound of the following formula:




- 5       , wherein  $R^1$  represents a cation, H, or  $C_{1-12}$ alkyl,  
 $R^2$  and  $R^3$  each represent H or a hydroxy protective group,  
 $R^4$  represents H or  $C_{1-3}$ alkyl, and  
 $R^5$  represents H or  $C_{1-6}$ alkyl, comprising the steps of:  
 (1) converting a compound of the following formula:



10


- , wherein  $R^{3a}$  represents a hydroxy protective group, and  
 $R^4$  and  $R^5$  are each as defined above,  
 into its cuprate, and then, performing stereo-specific 1,4-addition reaction of the  
 cuprate to an  $\alpha$ ,  $\beta$ -unsaturated ketone of the following formula:



15


- , wherein  $R^1$  is as defined above,  
 $R^{2a}$  represents a hydroxy protective group, and  
 $R'$  represents a hydroxy protective group,

to form a compound of the following formula;




, wherein  $R^1$ ,  $R^{2a}$ ,  $R^{3a}$ ,  $R^4$ ,  $R^5$  and  $R'$  are each as defined above;

(2) reducing a ketone of cyclopentanone in the compound of formula (IV) to form  
5 an  $\alpha$ -alcohol compound of the following formula:



, wherein  $R^1$ ,  $R^{2a}$ ,  $R^{3a}$ ,  $R^4$ ,  $R^5$ , and  $R'$  are each as defined above;


(3) substituting the  $\alpha$ -alcohol of the compound of formula (V) with a halide to form a  $\beta$ -halide compound of the following formula:



10

, wherein  $R^1$ ,  $R^{2a}$ ,  $R^{3a}$ ,  $R^4$ ,  $R^5$  and  $R'$  are each as defined above, and  
X represents halo;

(4) deprotecting a hydroxy protective group of the compound of formula (VI) to form a compound of the following formula:



(VII)

, wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ , and  $X$  are each as defined above;

(5) performing intramolecular cyclization to the compound of formula (VII) to form the compound of formula (I).

5

2. The process of Claim 1, wherein  $R^2$ ,  $R^3$ ,  $R^{2a}$  and  $R^{3a}$  each represent trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, t-butyl diphenylsilyl, phenyl dimethylsilyl, or tetrahydropyranyl, and  $R^4$  represents methyl, methoxymethyl, methoxyethyl, benzyloxymethyl, or p-benzyloxymethyl.

10

3. The process of Claim 1, wherein in step (2) the reduction is performed with using a metal hydride.

4. The process of Claim 3, wherein the metal hydride is selected from the group consisting of sodium borohydride ( $\text{NaBH}_4$ ), L-selectride, N-selectride and K-selectride.

15

5. The process of Claim 1, wherein in step (3), i) the  $\alpha$ -alcohol is converted into a leaving group, and then, the leaving group is converted into the  $\beta$ -halide via  $\text{S}_{\text{N}}2$  reaction with the halide nucleophile, or ii) the  $\alpha$ -alcohol is directly converted into the  $\beta$ -halide using trialkylphosphine and carbon tetrahalide.


20

6. The process of Claim 1, wherein in step (4) the hydroxy protective group is deprotected under an acidic condition.

7. The process of Claim 1, wherein in step (5) the cyclization reaction is performed in the presence of base.

25

8. A compound of the following formula:



, wherein  $\text{R}^{3a}$ ,  $\text{R}^4$  and  $\text{R}^5$  are each as defined in Claim 1.

Rec'd PCT/PTC 17 MAR 2005  
PC03017-YS

## PCT REQUEST

Original (for SUBMISSION) - printed on 18.09.2003 10:17:25 AM

0	For receiving Office use only	
0-1	International Application No.	
0-2	International Filing Date	
0-3	Name of receiving Office and "PCT International Application"	
0-4	Form - PCT/RO/101 PCT Request	
0-4-1	Prepared using	PCT-EASY Version 2.92 (updated 01.04.2003)
0-5	Petition The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty	
0-6	Receiving Office (specified by the applicant)	Korean Intellectual Property Office (RO/KR)
0-7	Applicant's or agent's file reference	PC03017-YS
I	Title of invention	PROCESS FOR PREPARING PROSTAGLANDIN DERIVATIVES AND STARTING MATERIALS FOR THE SAME
II	Applicant	
II-1	This person is:	applicant only
II-2	Applicant for	all designated States except US
II-4	Name	YONSUNG FINE CHEMICAL CO. LTD.
II-5	Address:	51-301, Science Technology Park Sungkyunkwan University, 300, Chonchon-dong, Changan-gu, Suwon 440-746 Kyonggi-do Republic of Korea
II-6	State of nationality	KR
II-7	State of residence	KR
II-8	Telephone No.	82-31-290-7961
II-9	Facsimile No.	82-31-290-7962
II-10	e-mail	uncleduly@hanmail.net

## PCT REQUEST

PC03017-YS

Original (for SUBMISSION) - printed on 18.09.2003 10:17:25 AM

III-1	Applicant and/or inventor	
III-1-1	This person is:	applicant and inventor
III-1-2	Applicant for	US only
III-1-4	Name (LAST, First)	KIM, Yong Hyun
III-1-5	Address:	73-73, Hwagok 5-dong Gangseo-gu 157-015 Seoul Republic of Korea
III-1-6	State of nationality	KR
III-1-7	State of residence	KR
III-2	Applicant and/or inventor	
III-2-1	This person is:	applicant and inventor
III-2-2	Applicant for	US only
III-2-4	Name (LAST, First)	LEE, Yiu Suk
III-2-5	Address:	Woosung Apt. #304-407 105, Sadang-dong, Dongjak-gu 156-090 Seoul Republic of Korea
III-2-6	State of nationality	KR
III-2-7	State of residence	KR
IV-1	Agent or common representative; or address for correspondence The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:	agent
IV-1-1	Name (LAST, First)	CHOI, Kyu Pal
IV-1-2	Address:	4th Floor, Halla Classic Building 824-11, Yeoksam-dong, Kangnam-gu 135-080 Seoul Republic of Korea
IV-1-3	Telephone No.	82-2-555-6888
IV-1-4	Facsimile No.	82-2-555-9888
IV-1-5	e-mail	hansung@hsip.co.kr

## PCT REQUEST

PC03017-YS

Original (for SUBMISSION) - printed on 18.09.2003 10:17:25 AM

<b>V</b>	<b>Designation of States</b>	
<b>V-1</b>	Regional Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	<p>AP: GH GM KE LS MW MZ SD SL SZ TZ UG ZM ZW and any other State which is a Contracting State of the Harare Protocol and of the PCT</p> <p>EA: AM AZ BY KG KZ MD RU TJ TM and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT</p> <p>EP: AT BE BG CH&amp;LI CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE SI SK TR and any other State which is a Contracting State of the European Patent Convention and of the PCT</p> <p>OA: BF BJ CF CG CI CM GA GN GQ GW ML MR NE SN TD TG and any other State which is a member State of OAPI and a Contracting State of the PCT</p>
<b>V-2</b>	National Patent (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	<p>AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH&amp;LI CN CO CR CU CZ DE DK DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NI NO NZ OM PG PH PL PT RO RU SC SD SE SG SK SL SY TJ TM TN TR TT TZ UA UG US UZ VC VN YU ZA ZM ZW</p>
<b>V-5</b>	<b>Precautionary Designation Statement</b> In addition to the designations made under items V-1, V-2 and V-3, the applicant also makes under Rule 4.9(b) all designations which would be permitted under the PCT except any designation(s) of the State(s) indicated under item V-6 below. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit.	
<b>V-6</b>	<b>Exclusion(s) from precautionary designations</b>	NONE
<b>VI-1</b>	<b>Priority claim of earlier national application</b>	
VI-1-1	Filing date	18 September 2002 (18.09.2002)
VI-1-2	Number	10-2002-0056836
VI-1-3	Country	KR
<b>VII-1</b>	<b>International Searching Authority Chosen</b>	Korean Intellectual Property Office (KIPO) (ISA/KR)

## PCT REQUEST

PC03017-YS

Original (for SUBMISSION) - printed on 18.09.2003 10:17:25 AM

<b>VIII</b>	<b>Declarations</b>	<b>Number of declarations</b>	
VIII-1	Declaration as to the identity of the inventor	-	
VIII-2	Declaration as to the applicant's entitlement, as at the international filing date, to apply for and be granted a patent	-	
VIII-3	Declaration as to the applicant's entitlement, as at the international filing date, to claim the priority of the earlier application	-	
VIII-4	Declaration of inventorship (only for the purposes of the designation of the United States of America)	-	
VIII-5	Declaration as to non-prejudicial disclosures or exceptions to lack of novelty	-	
<b>IX</b>	<b>Check list</b>	<b>number of sheets</b>	<b>electronic file(s) attached</b>
IX-1	Request (including declaration sheets)	5	-
IX-2	Description	15	-
IX-3	Claims	4	-
IX-4	Abstract	1	EZABST00.TXT
IX-5	Drawings	0	-
IX-7	TOTAL	25	
	<b>Accompanying items</b>	<b>paper document(s) attached</b>	<b>electronic file(s) attached</b>
IX-8	Fee calculation sheet	✓	-
IX-9	Original separate power of attorney	✓	-
IX-13	Priority document(s)	Item(s) VI-1	-
IX-17	PCT-EASY diskette	-	Diskette
IX-19	Figure of the drawings which should accompany the abstract		
IX-20	Language of filing of the international application	English	
X-1	Signature of applicant, agent or common representative		
X-1-1	Name (LAST, First)	CHOI, Kyu Pal	

## FOR RECEIVING OFFICE USE ONLY

10-1	Date of actual receipt of the purported international application	
10-2	Drawings:	
10-2-1	Received	
10-2-2	Not received	
10-3	Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application	
10-4	Date of timely receipt of the required corrections under PCT Article 11(2)	
10-5	International Searching Authority	ISA/KR



## PCT REQUEST

PC03017-YS

Original (for SUBMISSION) - printed on 18.09.2003 10:17:25 AM

10-6	Transmittal of search copy delayed until search fee is paid	
------	---	--

## FOR INTERNATIONAL BUREAU USE ONLY

11-1	Date of receipt of the record copy by the International Bureau	
------	--	--

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**